

PREMIUM PRO PLUS

For Use by Licensed Professionals Only

This pre-deadener was originally developed by a pharmacist familiar with our industries and has been used successfully for decades. This recent revision has improved its speed and depth of anesthesia. This cream contains 1% Tetracaine and 5% Lidocaine and with a pH of 7 and is *safe for eyeliner*.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Store in a cool dark place, *out of the reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and contact an eye care physician. **DO NOT SWALLOW.** If swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES	EYEBROWS 15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine .
	EYELINER 15- 20 minutes
	Fill the eyes with eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a Q-tip applicator across the lash line. Remove anesthetic with a cold wet cotton ball before beginning your procedure. Following the first pass, apply Maximum Zone 2 or New and Improved Blue Gel to control swelling and maintain anesthesia for two hours.
	LIPS 15-20 minutes
For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine . Reapply Superior Super Juice during the procedure to continue anesthesia, or Five-Star Vasocaine , New and Improved Blue Gel or Maximum Zone 2 to continue anesthesia <u>and</u> constrict blood vessels.	
BODY SKIN 90-120 minutes	
Prior to tattooing, laser, or other procedure, lightly exfoliate, massage into the skin and then apply a medium coating over the top. To prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia <u>and</u> constrict blood vessels.	
PIERCING 15- 20 minutes	
Apply to eyebrows, navel and nipples. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate. Gently massage into skin, then apply a medium coating on top. Cover with occlusive dressing. Wipe away immediately prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling.	

DERMAL SOURCE

2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.

Phone: +1 866-568-3223, Fax: 503-595-0293

www.dermalsource.com



PREMIUM PRO PLUS

For Use by Licensed Professionals Only

This pre-deadener was originally developed by a pharmacist familiar with our industries and has been used successfully for decades. This recent revision has improved its speed and depth of anesthesia. This cream contains 1% Tetracaine and 5% Lidocaine and with a pH of 7 and is *safe for eyeliner*.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Store in a cool dark place, *out of the reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and contact an eye care physician. **DO NOT SWALLOW.** If swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES	EYEBROWS 15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine .
	EYELINER 15- 20 minutes
	Fill the eyes with eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a Q-tip applicator across the lash line. Remove anesthetic with a cold wet cotton ball before beginning your procedure. Following the first pass, apply Maximum Zone 2 or New and Improved Blue Gel to control swelling and maintain anesthesia for two hours.
	LIPS 15-20 minutes
For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine . Reapply Superior Super Juice during the procedure to continue anesthesia, or Five-Star Vasocaine , New and Improved Blue Gel or Maximum Zone 2 to continue anesthesia <u>and</u> constrict blood vessels.	
BODY SKIN 90-120 minutes	
Prior to tattooing, laser, or other procedure, lightly exfoliate, massage into the skin and then apply a medium coating over the top. To prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia <u>and</u> constrict blood vessels.	
PIERCING 15- 20 minutes	
Apply to eyebrows, navel and nipples. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate. Gently massage into skin, then apply a medium coating on top. Cover with occlusive dressing. Wipe away immediately prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling.	

DERMAL SOURCE

2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.

Phone: +1 866-568-3223, Fax: 503-595-0293

www.dermalsource.com

