

MAXIMUM ZONE 1

For Use by Licensed Professionals Only

This 5% Lidocaine topical anesthetic has the advantage of a pH of 7, for rapid onset and safe use around the eyes, although care should be taken to avoid getting the product in the eyes. The cream base also makes this product self-occlusive. This product is for relief of discomfort due to anorectal disorders, or in preparation for other pain sensitive procedures such as lasers, electrolysis, tattooing or piercing.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children.*

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and contact an eye care physician. *DO NOT SWALLOW.* If accidentally swallowed, seek medical help immediately.

| | |
|--|--|
| SUGGESTED APPLICATION TIMES | EYEBROWS 15-20 minutes |
| | We recommend using Premium Pro Plus for this procedure. Use Superior Super Juice during the procedure to continue anesthesia, and either Maximum Zone 2 or New and Improved Blue Gel to continue anesthesia and constrict blood vessels. |
| | EYELINER 10- 15 minutes |
| | Fill the eyes with liquid eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a small applicator across the lash line. Remove anesthetic before beginning procedure. Following the first pass of tattooing, sparingly apply Maximum Zone 2 or New and Improved Blue Gel to control swelling and maintain anesthesia. |
| | LIPS 15-20 minutes |
| We suggest using Premium Pro Plus or Top Shelf Super T for this procedure. While some relief may be provided with topical anesthetics, for total anesthesia, you must provide a nerve block performed by a physician. <i>Do not use inside of the mouth.</i> Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels. | |
| BODY SKIN 90-120 minutes | |
| Prior to procedures, lightly exfoliate and massage Maximum Zone 1 into skin and then apply a medium coating over the top. Cover with an occlusive dressing. Thicker skin requires longer set-up time than thinner skin. Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels. | |
| PIERCING 15- 20 minutes | |
| We recommend Top Shelf Super T for this procedure. Avoid genital areas. Thicker skin requires longer pre-treating time. Massage Top Shelf Super T into skin, then apply a coating on top. Wipe away prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling. | |

DERMAL SOURCE, INC
2319 NE Glisan Street, Portland, OR 97232, U.S.A.
Phone: +1 866-568-3223, Fax: 503-595-0293
www.dermalsource.com



MAXIMUM ZONE 1

For Use by Licensed Professionals Only

This 5% Lidocaine topical anesthetic has the advantage of a pH of 7, for rapid onset and safe use around the eyes, although care should be taken to avoid getting the product in the eyes. The cream base also makes this product self-occlusive. This product is for relief of discomfort due to anorectal disorders, or in preparation for other pain sensitive procedures such as lasers, electrolysis, tattooing or piercing.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children.*

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline and contact an eye care physician. *DO NOT SWALLOW.* If accidentally swallowed, seek medical help immediately.

| | |
|--|--|
| SUGGESTED APPLICATION TIMES | EYEBROWS 15-20 minutes |
| | We recommend using Premium Pro Plus for this procedure. Use Superior Super Juice during the procedure to continue anesthesia, and either Maximum Zone 2 or New and Improved Blue Gel to continue anesthesia and constrict blood vessels. |
| | EYELINER 10- 15 minutes |
| | Fill the eyes with liquid eye gel to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a small applicator across the lash line. Remove anesthetic before beginning procedure. Following the first pass of tattooing, sparingly apply Maximum Zone 2 or New and Improved Blue Gel to control swelling and maintain anesthesia. |
| | LIPS 15-20 minutes |
| We suggest using Premium Pro Plus or Top Shelf Super T for this procedure. While some relief may be provided with topical anesthetics, for total anesthesia, you must provide a nerve block performed by a physician. <i>Do not use inside of the mouth.</i> Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels. | |
| BODY SKIN 90-120 minutes | |
| Prior to procedures, lightly exfoliate and massage Maximum Zone 1 into skin and then apply a medium coating over the top. Cover with an occlusive dressing. Thicker skin requires longer set-up time than thinner skin. Use Superior Super Juice during procedures to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels. | |
| PIERCING 15- 20 minutes | |
| We recommend Top Shelf Super T for this procedure. Avoid genital areas. Thicker skin requires longer pre-treating time. Massage Top Shelf Super T into skin, then apply a coating on top. Wipe away prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling. | |

DERMAL SOURCE, INC
2319 NE Glisan Street, Portland, OR 97232, U.S.A.
Phone: +1 866-568-3223, Fax: 503-595-0293
www.dermalsource.com

