

HOW TO MAXIMIZE DERMAL SOURCE TOPICAL ANESTHETICS FOR TATTOOING, PERMANENT MAKEUP, LASER, PIERCING, ELECTROLYSIS AND OTHER PAINFUL PROCEDURES

Integrating topical anesthetics into your practice is easy to do once you understand how each product functions.

PRE-PROCEDURE ANESTHETICS FOR INTACT SKIN

To enhance absorption for most procedures, excluding eyeliner: 1) exfoliate and clean the skin with an alcohol pad; 2) roll the area with a Derm Ease G2 roller; 3) apply anesthetic; and 4) occlude until the anesthetic effect is achieved.

TOP SHELF SUPER T: This triple anesthetic predeadener cream is effective on thicker skin applications, body piercing, electrolysis, laser treatments and other painful procedures. It has a pH of 5.5, which allows for penetration of the epidermis without irritation.

PREMIUM PRO PLUS: This self-occlusive predeadener cream contains the highest percentage of Lidocaine and Tetracaine available without a prescription. With a pH of 7, it is safe for use on all procedures including eyeliner.

MAXIMUM ZONE 1: This self-occlusive predeadener cream contains the highest percentage of Lidocaine available without a prescription, and with a pH of 7, it is safe for use on all procedures including eyeliner.

TOPICAL ANESTHETICS FOR USE DURING PAINFUL PROCEDURES

SUPERIOR SUPER JUICE: This liquid anesthetic contains Benzocaine, and the highest levels of Tetracaine and Lidocaine available without a prescription. Apply it to the broken skin as you work and receive an anesthetic effect in 90 seconds. Excluding eyeliner, use it in place of soap or alcohol, which stop the action of your anesthetics and change the pH of your skin. This product can be used to remove your pre-procedure topical anesthetics, excluding eyeliner. Unlike **New and Improved Blue Gel** and **Maximum Zone 2**, this product cannot be overused on the skin.

NEW AND IMPROVED BLUE GEL: This gel anesthetic contains the highest level of Tetracaine and Lidocaine available without a prescription, together with a vascular constrictor to reduce swelling and bleeding.

MAXIMUM ZONE 2: This clear gel anesthetic contains the highest level of Lidocaine available without a prescription, together with a vascular constrictor to reduce swelling and bleeding.

NEW AND IMPROVED BLUE GEL AND MAXIMUM ZONE 2 can be occluded and are effective for up to two hours with one application. They are safe for use on all procedures. Thoroughly remove them before you begin your work. Do not reapply.

Apply **Five-Star Vasocaine** or **Superior Super Juice** to extend the anesthetic effect, excluding eyeliner.

FIVE-STAR VASOCAINE: This liquid spray with a vascular constrictor contains the highest level of Lidocaine available without a prescription. It is safe for use in all procedures, except eyeliner. Spray it on open skin. Because of the vascular constrictor, you will also experience a reduction in bruising and bleeding. Always remove the product thoroughly before continuing a procedure.

ADDITIONAL TOPICAL ANESTHETIC INFORMATION

1. Keep your anesthetics refrigerated or in a cool dark place. Do not freeze your anesthetics. Temperatures above 80 degrees reduce the life and effectiveness of your anesthetics. Do not use beyond expiration date.
2. Remove your pre-procedure anesthetics before applying a design, stencil or landmarks for your procedure.
3. For body piercing, dip your penetrator bar into **New and Improved Blue Gel** or **Maximum Zone 2** to reduce swelling and bleeding, and aid in the insertion of the jewelry.

DERMAL SOURCE, INC. Phone: +1 866-568-3223, Fax: 503-595-0293
2319 NE Glisan Street, Portland, Oregon 97232, U.S.A. www.dermalsource.com

04/13/2021





	Pre-Procedure	During Procedure	Long Lasting	Fast Setup	Reduces Swelling	Eyeliner	Lips	Eyebrow	Tattoos	Piercing	Laser/Dermabrasion	Electrology
Premium Pro Plus (cream) 1% tetracaine 5% lidocaine	X					X	X	X	X			
Top Shelf Super T (cream) 1% tetracaine 5% lidocaine 12% benzocaine	X						X	X	X	X	X	X
Maximum Zone 1 (cream) 5% lidocaine	X					X		X	X			
Maximum Zone 2 (gel) 5% lidocaine 0.1% epinephrine		X	X		X	X	X	X	X	X	X	X
New and Improved Blue Gel 1% tetracaine 5% lidocaine 0.1% epinephrine		X	X		X	X	X	X	X	X	X	X
Superior Super Juice (liquid) 1% tetracaine 5% lidocaine 12% benzocaine		X		X			X	X	X			
Five-Star Vasocaine (liquid spray) 5% lidocaine 0.1% epinephrine		X	X	X	X		X	X	X	X		